

Selecting native entomopathogenic fungi against *Cosmopolites sordidus* (Germar) in the laboratory

Seleção de fungos entomopatogênicos nativos contra Cosmopolites sordidus (Germar) em laboratório

Junior Steve Garrido Pérez^{1,2}, Richard Paredes-Espinosa³, Giannfranco Egoávil Jump¹, Oniel Jeremias Aguirre Gil^{3,4*}

¹Universidad Nacional Agraria de la Selva, Tingo María, Peru.

²Palmas del Espino S.A., Uchiza, Perú.

³Instituto Nacional de Innovación Agraria, Tarapoto, Peru.

⁴Universidade Estadual Paulista "Júlio de Mesquita Filho", Jaboticabal, SP, Brasil. * Author for correspondence: oniel.aguirre@unesp.br

Submission: 27/07/2020 | Acceptance: 14/12/2020

ABSTRACT

The objective of this study was to select native isolates of entomopathogenic fungi against adult *Cosmopolites sordidus* in laboratory conditions. Four isolates of *Beauveria bassiana*, two isolates of *Metarhizium anisopliae*, and two isolates of *Paecilomyces lilacinus* were tested against *C. sordidus*. The entomopathogenic fungi that resulted in mortality rates over 50% were *M. anisopliae* (MA-CsCha and MA-Carabid) and *B. bassiana* (BB-CsCha and BB-CsLp). The most virulent isolates were MA-CsCha, BB-CsCha, and MA-Carabid with median lethal times (LT50) of 4.82, 5.4, and 5.79 days, respectively. In conclusion, the MA-CsCha, BB-CsCha, MA-Carabid, and BB-CsLp isolates are viable candidates to be tested in banana fields.

KEYWORDS: banana root borer, *Beauveria bassiana*, *Metarhizium anisopliae*, biocontrol.

RESUMO

Este estudo teve como objetivo selecionar isolados de fungos entomopatogênicos contra adultos de *Cosmopolites sordidus* em condições de laboratório. Quatro isolados de *Beauveria bassiana*, dois isolados de *Metarhizium anisopliae* e dois isolados de *Paecilomyces lilacinus* foram testados contra *C. sordidus*. Os fungos entomopatogênicos que resultaram em mais de 50% da mortalidade de adultos de *C. sordidus* foram *M. anisopliae* (MA-CsCha e MA-Carabid) e *B. bassiana* (BB-CsCha e BB-CsLp). Os isolados mais virulentos foram MA-CsCha, BB-CsCha e MA-Carabid com um tempo letal médio (TL50) de 4,82, 5,4 e 5,79 dias, respectivamente. Em conclusão, os isolados MA-CsCha, BB-CsCha, MA-Carabid e BB-CsLp são candidatos para serem testados em plantações de banana.

PALAVRAS-CHAVE: Moleque-da-bananeira, *Beauveria bassiana*, *Metarhizium anisopliae*, biocontrole.

The banana crop is important because it contributes to food security with an annual production of about 127 million tons worldwide (NYINE & PILLAY 2007, FAOSTAT 2020). However, the commercial crop faces problems caused by pests, and the Banana Root Borer (BRB), *Cosmopolites sordidus* (Germar), which infests the cortical tissues, can generate losses of up to 100% of the harvest (AKELLO et al. 2008, OLIVEIRA et al. 2017).

In this context, entomopathogenic fungi are an integrated pest management tool and an important alternative to insecticides against the BRB (OLSON 2015). There is a wide distribution of entomopathogenic fungi, which means they have a high genetic variability and distinct levels of virulence and local adaptability (NIASSY et al. 2013). Therefore, suitable isolates should be selected to control BRB populations according to the abovementioned aspects, since these studies are scarce under Peruvian conditions.

The entomopathogens *Beauveria bassiana* (Bals.) Vuill. (now *Cordyceps bassiana*) and *Metarhizium anisopliae* (Metsch.) Sorokin have been shown to be the most promising biocontrol agents against BRB due to their cosmopolitan distribution and high lethality caused by their production of mycotoxins and growth which blocks the insect's digestive tract (TINZAARA et al. 2015, SALUSTINO et al. 2019). These species are

reported to cause infections for more than 300 species of pest insects (MOREIRA et al. 2017), which implies that this high genetic variability should be studied in order to select efficient entomopathogenic fungi adapted to the conditions where it is necessary to control BRB. Furthermore, the discovery of new native isolates with high virulence and infectivity can add value to biological control methods by reducing production costs and the use of conventional insecticides in banana fields (MEMBANG et al. 2020).

The objective of this study was to select native entomopathogenic fungi as potential biocontrol agents of *C. sordidus* populations in a humid tropic climate. The hypothesis was that at least one entomopathogenic fungus in the study would reach a 50% mortality rate within seven days after inoculation. Therefore, the results of this study offer new opportunities to discover potential entomopathogenic fungi that can be used in field using local and exotic isolates.

The study was conducted in the Laboratory of Entomopathogens at the Universidad Nacional Agraria de la Selva (UNAS), Tingo María, Peru (9°18'48" S, 75°59'45" W, 668.6 m above sea level), which has a humid tropical climate (Af) based on the Köppen-Geiger classification (PEEL et al. 2007). The eight entomopathogenic fungi tested in this study were obtained from the Servicio Nacional de Sanidad Agraria (SENASA) and UNAS collections. In the laboratory, four isolates of *Beauveria bassiana*, two isolates of *Metarhizium anisopliae*, and two isolates of *Paecilomyces lilacinus* (Tom) Samson were tested against *C. sordidus*. Information about the collections is presented in Table 1.

The entomopathogenic fungi were cultured in Petri dishes (8.5 cm diameter) containing Potato Dextrose Agar (PDA) at 24±5 °C for ten days. The Petri dishes and PDA were previously autoclaved at 121 °C, 15 lbs for 15 minutes.

After ten days, entomopathogenic fungi were sown in an artificial substrate. The artificial substrate had been prepared as follows: 200 g rice and 40 mL water placed in a polypropylene bag (8 x 12 cm); the bags were then folded, stapled closed, and autoclaved at 121 °C, 15 lbs for 15 minutes. Entomopathogenic fungi were sown in an artificial substrate in a laminar flow cabinet.

The bags with entomopathogenic fungi were stored on metal shelves in a room at 24 ± 5 °C and incubated for ten days. The bags were shaken daily to homogenize the internal content.

Table 1. Description of the isolates.

Isolate codes	Scientific name	Origin
BB-Hh	<i>Beauveria bassiana</i>	<i>Hypothenemus hampei</i> collected in Sania Province, Peru (SENASA, commercial code: CCB-LE265)
BB-CsCha	<i>Beauveria bassiana</i>	<i>Cosmopolites sordidus</i> collected in Chanchamayo Province, Peru (SENASA, commercial code: CCB-LE2118)
BB-CsLp	<i>Beauveria bassiana</i>	<i>Cosmopolites sordidus</i> collected in Leoncio Prado Province, Peru (UNAS)
BB-CsVr	<i>Beauveria bassiana</i>	<i>Cosmopolites sordidus</i> collected in Villa Rica District, Peru (UNAS)
PL-Milt	<i>Paecilomyces lilacinus</i>	<i>Meloidogyne incognita</i> collected in Italy (SENASA, commercial code: CCB-LE 701)
PL-CsChi	<i>Paecilomyces lilacinus</i>	<i>Cosmopolites sordidus</i> collected in Chiclayo Province (UNAS)
MA-Carabid	<i>Metarhizium anisopliae</i>	Unidentified Carabid specimen in Peru (SENASA, commercial code: CCB-LE302)
MA-CsCha	<i>Metarhizium anisopliae</i>	<i>Cosmopolites sordidus</i> collected in Chanchamayo Province, Peru (SENASA, commercial code: CCB-LE319)
Control	-	Sterile water

For the conidial suspension, a ten-day-old fungal colony was diluted in sterile distilled water and mixed with Tween 80[®] (0.05%). Then, the mixture was filtered through sterilized gauze to separate the conidia from the mycelium and other impurities. The quantity of conidia contained in the mixture were registered using a Neubauer chamber and then adjusted to a concentration of 10⁸ conidia mL⁻¹. This concentration was used for the findings obtained by MEMBANG et al. (2020).

Adult *Cosmopolites sordidus* were collected in a banana field at Los Milagros Village, Tingo María, Peru. Traps formed by 10 cm banana stalks were used to attract *C. sordidus* in the field. One week later, adult *C. sordidus* were collected from the traps and placed in plastic boxes to be transported. In the laboratory, the adult *C. sordidus* were sprayed with and immersed in sodium hypochlorite (0.5%) for disinfection and then rinsed with sterile water three times. Only the more active ones of these insects were

used in the experiment.

A completely randomized design with nine treatments was adopted to determine the mortality rates and lethal times of entomopathogenic fungi isolates (Table 1) against *C. sordidus* in the laboratory.

For each isolate, an experiment was conducted with three replicates. One replicate consisted of a group of 15 healthy adult *C. sordidus* that were individually sprayed with the solution containing the entomopathogens until they were dripping. A total of 45 adult *C. sordidus* were used per isolate. The sprayed insects were then conditioned in Petri dishes (10 cm diameter) containing banana stalks, which had been disinfected with 0.5% sodium hypochlorite solution for five min, in order to feed.

The number of active *C. sordidus* were evaluated for 21 days after infection. The mortality rate was calculated by the percentage of dead insects from total insects. Dead insects were individually maintained in Petri dishes lined with moistened filter paper to allow fungal sporulation.

The mortality rate percentage (x) under laboratory conditions was transformed by arcsine $(x/100)^{1/2}$. To obtain the median lethal times (LT50) of the isolates against *C. sordidus*, a Probit analysis was conducted. Data were analyzed using the R software for Windows (R CORE TEAM 2016). Cumulative mortality rates were corrected using Abbott's formula. Normality and equality of variance were tested by the Kolmogorov-Smirnov and the Levene's tests, respectively ($p \leq 0.05$). The mortality rate percentage was analyzed as a response variable in one-way (isolate) ANOVA. Whenever a significant F-test ($p \leq 0.05$) was detected, the mean values for mortality were compared with Tukey's test (HSD) at ($p \leq 0.05$). The values of LT50 and LT90 of the isolates were estimated at 95% confidence levels (CL) using the Probit analysis with the "ecotoxicology" Package (GAMA 2015).

In total, 878 adult *C. sordidus* were collected for the study. The experiment revealed the ability of entomopathogenic fungi isolates to control *C. sordidus* adults under laboratory conditions. The MA-CsCha isolate resulted in a 57.78% mortality rate for adult *C. sordidus* five days after inoculation in the laboratory (Table 2). However, the MA-CsCha, BB-CsCha, MA-Carabid, and BB-CsLp isolates resulted in mortality rates of over 50% for adult *C. sordidus* seven days after inoculation.

According to the Probit regression, the MA-CsCha isolate resulted in a 50% mortality rate for *C. sordidus* 4.82 days after inoculation (Figure 1A). However, the BB-CsCha isolate resulted in a 90% mortality rate for *C. sordidus* in 8.04 days (Figure 1C). The differences between isolates are explained by the slope indicating that the rate of mortality increases 7.44 times per day for adult *C. sordidus* inoculated with BB-CsCha.

Table 2. Mortality rates (%) for adult *Cosmopolites sordidus* infected with entomopathogenic fungi in the laboratory.

Treatments	Days after infection		
	5	6	7
MA-CsCha	57.78a	71.11a	82.22a
BB-CsCha	24.44b	77.78a	82.22ab
MA-Carabid	8.89b	73.33a	84.44a
BB-Hh	13.33b	28.89b	33.33cd
BB-CsLp	13.33b	24.44bc	53.33bc
PL-Milt	4.44b	4.44c	4.44e
PL-CsChi	4.44b	6.67bc	11.11de
BB-CsVr	2.22b	6.67bc	11.11de

Mean values followed by different letters in the same column are significantly different based on Tukey's test ($p \leq 0.05$)

Metarhizium anisopliae and *B. bassiana* infect a wide range of hosts in numerous insect orders, including Orthoptera, Lepidoptera, Dermaptera, Diptera, and Coleoptera (MAINA et al. 2018). However, in order to select isolates and use them for biological control, it is fundamental to understand the mortality rate and lethal time (kill at least 50% individuals) of an entomopathogenic fungus on a specific target pest (BAYISSA et al. 2017, MWEKE et al. 2018, MAHOT et al. 2019). This study reports on two isolates of *M. anisopliae* (MA-CsCha, MA-Carabid) and two isolates of *B. bassiana* (BB-CsCha, BB-CsLp), which can cause mortality rates of over 50% for adult *C. sordidus* under laboratory conditions. In Brazil, FANCELLI et al. (2013) reported similar results, wherein native isolates of *B. bassiana* promoted mortality rates (14 - 96%) for adult *C. sordidus* in laboratory experiments. The most virulent isolates achieved a LT50 of 4.82, 5.40, and 5.79 days (Figure 1), which were faster than the most virulent isolates reported by FANCELLI et al. (2013),

GONZÁLEZ et al. (2018), and LOPES et al. (2011).

On the other hand, some studies have reported that exotic isolates can possibly be more effective than native isolates (LOPES et al. 2013). Although it is not a rule, this study used exotic isolates from Sania (BB-Hb), Villa Rica (BB-CsVr), Chiclayo Province (PL-CsChi), and Italy (PL-Milt), which did not present relevant results.




Figure 1. Mortality rates for adult *Cosmopolites sordidus* caused by infection from isolates MA-CsCh (A), BB-CsCh (B), and MA-Carabid (C) over time, according to the Probit regression.

The findings also indicate variations between and within entomopathogenic fungi isolates, as reported by CHENG et al. (2016). Therefore, an individual belonging to the same fungal species is not a guarantee for similar virulence, because isolates can use different types and quantities of enzymes and metabolites to infect the insect host (ABDELAZIZ et al. 2018).

Future studies should use combinations of different species of biological control agents so as to increase the effects against pests, reduce insecticide use, and minimize environmental pollution risks and pest resistance (REDDY et al. 2014, CARVALHO 2017).

The results of this study indicate suitable entomopathogenic candidates to be tested in banana crops against *C. sordidus*. In conclusion, the native isolates of *Metarhizium anisopliae* (MA-CsCha and MA-Carabid) and *Beauveria bassiana* (BB-CsCha and BB-CsLp) were selected as potential biocontrol agents against *C. sordidus*.

ACKNOWLEDGEMENTS

The authors thank the Facultad de Agronomía - Universidad Nacional Agraria de la Selva and the Estación Experimental Agraria El Porvenir - Instituto Nacional de Innovación Agraria for their support in this project.

REFERENCES

- ABDELAZIZ O et al. 2018. Pathogenicity of three entomopathogenic fungi, to the aphid species, *Metopolophium dirhodum* (Walker) (Hemiptera: Aphididae), and their Alkaline protease activities. *Egyptian Journal of Biological Pest Control* 28: 24.
- AKELLO J et al. 2008. Effect of endophytic *Beauveria bassiana* on populations of the banana weevil, *Cosmopolites sordidus*, and their damage in tissue-cultured banana plants. *Entomologia Experimentalis et Applicata* 129: 157-165.
- BAYISSA W et al. 2017. Selection of fungal isolates for virulence against three aphid pest species of crucifers and okra. *Journal of Pest Science* 90: 355-368.
- CARVALHO FP. 2017. Pesticides, environment, and food safety. *Food and Energy Security* 6: 48-60.
- CHENG Y et al. 2016. Evaluation of Pathogenicity of the fungi *Metarhizium anisopliae* and *Beauveria bassiana* in hazelnut weevil (*Curculio nucum* L., Coleoptera, Curculionidae) Larvae. *Indian Journal of Microbiology* 56: 405-410.
- FANCELLI M et al. 2013. *Beauveria bassiana* strains for biological control of *Cosmopolites sordidus* (Germ.) (Coleoptera: Curculionidae) in plantain. *BioMed Research International* 2013: 1-7.
- FAOSTAT. 2020. Crop statistics. Database. Available in: <http://www.fao.org/faostat/en/#data/QC>. Accessed in: 27 jul. 2020.
- GAMA J. 2015. Ecotoxicology: Methods for Ecotoxicology (R package version 1.0.1) [Computer software]. Available in: <https://CRAN.R-project.org/package=ecotoxicology>. Accessed in: 27 feb. 2020.
- GONZÁLEZ DN et al. 2018. Suitability of *Cordyceps bassiana* and *Metarhizium anisopliae* for biological control of *Cosmopolites sordidus* (Germar) (Coleoptera: Curculionidae) in an organic Mexican banana plantation: laboratory and field trials. *Journal of Plant Diseases and Protection* 125: 73-81.
- LOPES RB et al. 2013. Diversity of indigenous *Beauveria* and *Metarhizium* spp. in a commercial banana field and their virulence toward *Cosmopolites sordidus* (Coleoptera: Curculionidae). *Fungal Ecology* 6: 356-364.
- LOPES RB et al. 2011. Virulence and horizontal transmission of selected Brazilian strains of *Beauveria bassiana* against *Cosmopolites sordidus* under laboratory conditions. *Bulletin of Insectology* 64: 201-208.
- MAHOT HC et al. 2019. Laboratory assessment of virulence of Cameroonian isolates of *Beauveria bassiana* and *Metarhizium anisopliae* against mirid bugs *Sahlbergella singularis* Haglund (Hemiptera: Miridae). *African Entomology* 27: 86-96.
- MAINA UM et al. 2018. A review on the use of entomopathogenic fungi in the management of insect pests of field crops. *Journal of Entomology and Zoology Studies* 6: 27-32.
- MEMBANG G et al. 2020. *Cosmopolites sordidus* (Germar) susceptibility to indigenous Cameroonian *Beauveria bassiana* (Bals.) Vuill. and *Metarhizium anisopliae* (Metsch.) isolates. *Journal of Applied Entomology* 144: 468-480.
- MOREIRA FJC et al. 2017. Controle de *Cosmopolites sordidus* (Coleoptera: Curculionidae) com os fungos entomopatogênicos *Beauveria bassiana* e *Metarhizium anisopliae* em banana. *Revista Verde de Agroecologia e Desenvolvimento Sustentável* 12: 366-373.
- MWEKE A et al. 2018. Evaluation of the entomopathogenic fungi *Metarhizium anisopliae*, *Beauveria bassiana* and *Isaria* sp. for the management of *Aphis craccivora* (Hemiptera: Aphididae). *Journal of Economic Entomology* 111: 1587-1594.
- NIASSY S et al. 2013. Use of *Metarhizium anisopliae* Chitinase genes for genotyping and virulence characterization. *BioMed Research International* 2013: 1-9.
- NYINE M & PILLAY M. 2007. Banana nectar as a medium for testing pollen viability and germination in *Musa*. *African Journal of Biotechnology* 6: 1175-1180.
- OLIVEIRA F et al. 2017. Infestation of the banana root borer among different banana plant genotypes. *Ciência Rural* 47: e20151114.
- OLSON S. 2015. An analysis of the biopesticide market now and where it is going. *Outlooks on Pest Management* 26: 203-206.
- PEEL MC et al. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology Earth System Scientia* 11: 1633-1644.
- R CORE TEAM. 2016. R: A Language and Environment for Statistical Computing (3.3.1) [En]. R Foundation for Statistical Computing. Available in: <https://www.R-project.org/>. Accessed in: 27 jul. 2020.
- REDDY GVP et al. 2014. Laboratory and field efficacy of entomopathogenic fungi for the management of the sweetpotato weevil, *Cylas formicarius* (Coleoptera: Brentidae). *Journal of Invertebrate Pathology* 122: 10-15.
- SALUSTINO ADS et al. 2019. Olfactory attraction of rhizome borer (Coleoptera: Curculionidae) to banana genotypes inoculated with entomopathogenic fungus. *Journal of Experimental Agriculture International* 40: 1-6.
- TINZAARA W et al. 2015. Enhancing dissemination of *Beauveria bassiana* with host plant base incision trap for the management of the banana weevil *Cosmopolites sordidus*. *African Journal of Agricultural Research* 10: 3878-3884.